

Biotechnology Journal International

Volume 29, Issue 1, Page 1-10, 2025; Article no.BJI.128488 ISSN: 2456-7051 (Past name: British Biotechnology Journal, Past ISSN: 2231–2927, NLM ID: 101616695)

# Pathogenic Susceptibility of Cashew (Anacardium occidentale L.) to Twelve Isolates of Colletotrichum sp Present on Six Weeds in Cashew Orchards in Côte d'Ivoire

# Traoré Aboulaye <sup>a\*</sup>, Soro Sibirina <sup>a</sup>, Ayémou Alloua R. E. <sup>a</sup>, Soro N. Arsène <sup>b</sup>, Traoré-Ouattara Karidia <sup>a</sup> and Koné Daouda <sup>b</sup>

 <sup>a</sup> Laboratoire d'Amélioration de la Production Agricole, UFR Agroforesterie, Université Jean Lorougnon Guédé, Daloa, Côte d'Ivoire.
<sup>b</sup> African Centre of Excellence on Climate Change, Biodiversity and Sustainable Agriculture (CEA-CCBAD), Université Félix Houphouët-Boigny (UFHB), Abidjan, Côte d'Ivoire.

# Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/bji/2025/v29i1760

**Open Peer Review History:** 

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/128488

> Received: 17/10/2024 Accepted: 20/12/2024 Published: 01/01/2025

**Original Research Article** 

\*Corresponding author: E-mail: traoreaboulaye044@gmail.com;

**Cite as:** Aboulaye, Traoré, Soro Sibirina, Ayémou Alloua R. E., Soro N. Arsène, Traoré-Ouattara Karidia, and Koné Daouda. 2025. "Pathogenic Susceptibility of Cashew (Anacardium Occidentale L.) to Twelve Isolates of Colletotrichum Sp Present on Six Weeds in Cashew Orchards in Côte d'Ivoire". Biotechnology Journal International 29 (1):1-10. https://doi.org/10.9734/bji/2025/v29i1760.

# ABSTRACT

Cashew nuts are one of the agricultural products that contribute significantly to Côte d'Ivoire's economic development. However, cashew nut production in Côte d'Ivoire is threatened by anthracnose. Reducing anthracnose proliferation factors could reduce its impact on cashew trees. The aim of the study was to assess the virulence of *Colletotrichum* sp. isolated from weeds in cashew trees. It assessed the pathogenicity of 12 isolates of Colletotrichum sp. extracted from six weeds on cashew plants. The inocula were prepared by scraping the mycelium of each fungus in 10 ml of sterile distilled water. The spore suspension was collected and calibrated at 4.6.106 conidia/ml using a Malassez cell. This suspension was applied by spraying to the leaves of 30-dayold cashew plants. The parameters assessed were symptom incidence and severity index. All isolates tested induced symptoms characteristic of anthracnose on cashew plants. There was a significant difference between isolates in terms of their virulence. The highest incidence of 90% was obtained on plants inoculated with the CoIE2 isolate and the lowest incidence of 30% was obtained on plants inoculated with the ColN3 isolate. The highest severity index of 7.66 was obtained on plants inoculated with the CoIE3 isolate and the lowest severity index of 2.00 was obtained on plants inoculated with the CoIN3 isolate. These results show that Daniellia oliveri (Rolfe), Vitellaria paradoxa (G), Micuna pruriens (L), Blighia sapida (K. Koenig), Albizia zygia (DC) and Pterocarpus erinaceus (Poir) are hosts of anthracnose in cashew orchards in Côte d'Ivoire. Their integration into cashew pathogen control strategies is necessary.

Keywords: Inoculation; anthracnose; incidence; pathogenicity; Côte d'Ivoire.

# 1. INTRODUCTION

The cashew tree (Anacardium occidentale L.) is a perennial plant cultivated mainly for its nuts cashew Belonging and apples. to the Anacardiaceae family, the cashew tree has a high commercial value due to the food and industrial value of the cashew kernel. It is the only species of the Anacardium genus that is of economic interest because of its ability to produce nutritious apples and nuts (Agostini-Costa et al., 2006). As with other crops, abiotic and biotic factors constitute constraints to cashew cultivation. Among the biotic constraints, anthracnose is the major fungal disease of this crop in several producing countries (Wonni et al., 2017; Muntala et al., 2020; Veloso et al., 2022). This disease has a significant impact on the potential production of cashew trees. Anthracnose attacks leaves, inflorescences, nuts, and cashew apples, resulting in huge yield losses of more than 50% (Houndahouan et al., 2018). This disease in cashew is caused by the Colletotrichum gloeosporioides species complex including Colletotrichum chrysophilum, Colletotrichum fragariae, Colletotrichum fructicola, Colletotrichum gloeosporioides sensu Colletotrichum queenslandicum, stricto. Colletotrichum siamense and Colletotrichum tropicale (Veloso et al., 2018). In Côte d'Ivoire, anthracnose is present on cashew trees in all orchards, with varving degrees of severity from one site to another (Soro et al., 2015; Soro et al., 2022). It manifests itself through typical

symptoms such as patches of necrosis in the form of leaf scorch, scattered black dots on leaf blades, black spots on inflorescences and necrosis on cashew apples (Soro et al., 2013; Silué et al., 2017; Soro et al., 2022). In addition to cashew, the C. gloeosporioides species complex attacks other fruit plants (Figueiredo et The presence of alternative al.. 2012). Colletotrichum host plants in a field increases the risk of anthracnose spreading to cultivated plants (Dentika et al., 2021). The work of (Aboulaye et al., 2023) showed the presence of Colletotrichum fungi on weeds in the Ivorian cashew orchard. However, the pathogenic nature of these fungi on cashew remains to be demonstrated. The aim of this study was to assess the virulence of Colletotrichum sp. isolated from weeds in cashew crops. Specifically, the aim was to determine the latency period for the effect of the isolates on cashew plants and then to assess their incidence and the severity of the symptoms induced in a controlled environment.

# 2. MATERIALS AND METHODS

# 2.1 Study Area

The study was carried out at the Agricultural Production Improvement Laboratory of the Jean Lorougnon Guédé University. This university is located in the town of Daloa in central-western Côte d'Ivoire. The town of Daloa is located at latitude 6°53 north and longitude 6°27 west. Daloa is 139.1 km from Yamoussoukro, the political capital of Côte d'Ivoire, via the A6 motorway. It is the capital of the Haut-Sassandra Region, with a surface area of 15,200 km2 and an estimated population of more than 1,396977 (RGPH, 2021). The climate is Sudano-Guinean, with four seasons. The main rainy season lasts from April to mid-July, while the short dry season lasts from mid-July to mid-September. The short rainy season runs from mid-September to mid-November and the long dry season from December to March. The dry and wet seasons alternate, with temperatures ranging from 24.65°C to 27.75°C on average. Daloa's soil substrate is made up of old Precambrian granite. The region's soils are predominantly ferralitic. They are generally very deep, with high levels of organic matter. They have good agricultural potential for all types of crop (Zro et al., 2016).

#### 2.2 Materials

This work required the use of a set of materials comprising plant material, technical material and fungal material. The plant material consisted of 30-day-old cashew seedlings. These plants were produced on sterile substrate in nursery bags and pots. The substrate consisted of 50% soil and 50% wood soot. The substrate was steam sterilised for 60 minutes in a hermetically sealed metal bucket. The equipment used included an autoclave, a laminar flow hood, a Malassez cell, an electronic balance and an electron microscope. PDA (Potato Dextrose Agar) culture

medium was used to grow the fungi. The fungal material consisted of 12 isolates of Colletotrichum sp present on weeds in the cashew orchard (Table 1).

#### 2.3 Methods

#### 2.3.1 Study design

The system used was a completely randomised block with 13 treatments and 3 replicates. Each *Colletotrichum* sp isolate corresponded to a treatment that was applied to 05 cashew plants for one replication. Control plants were treated with tap water.

#### 2.3.2 Preparation of fungal inocula

The inoculum was prepared according to the method of (Silué et al., 2018), which was modified and adapted to the conditions of our study. The pure fungal cultures used for inoculum production were aged for 15 days on PDA medium. Preparation consisted of scraping the mycelium from each mushroom in 10 ml of sterile distilled water using a sterile scalpel. The spore suspension was collected using a sterile 20 ml syringe and filtered using filter paper to mycelium. remove any residual The concentration of conidia was calibrated at 4.6.106 conidia/ml using the Malassez cell. A drop of tween 20 was added to the spore suspension to facilitate its adhesion to the inoculated leaves.



Fig. 1. Map of the study area (Kouadio, 2015)

Fungal Genus	Code	Host Weed Zone	Agro-ecological
Colletotrichum	ColN1	Daniellia oliveri	North
	CoIN2	Pterocarpus erinaceus	North
	CoIN3	Vitellaria paradoxa	Nord
	CoIC1	Daniellia oliveri	Centre
	CoIC2	Blighia sapida	Centre
	CoIC3	Vitellaria paradoxa	Centre
	CoIE1	Daniellia oliveri	East
	CoIE2	Vitellaria paradoxa	East
	CoIE3	Albizia zygia	East
	ColCO1	Blighia sapida	Central West
	ColCo2	Pterocarpus erinaceus	Central West
	ColCo3	Micuna pruriens	Central West

Table 1. Fungal isolates used for I	pathogenicity tes	ting on cashew seedlings
	·····	

Grade	Surface area of infected	Interpretation
0	0 %	No symptoms
1	1-5 %	Moyenne
3	6-10 %	Medium
5	11-25 %	Mildly severe
7	26-50 %	Severe
9	>50 %	Very severe

# 2.3.3 Application of inoculum to cashew seedlings

The inoculum was applied to 30-day-old cashew plants. One day before inoculation, the plants were covered with transparent plastic to maintain humidity. The virulence of each fungus was tested on 05 healthy cashew seedlings. The inoculum was applied to the leaves of each cashew plant. After inoculation, the plants were covered with transparent plastic, depending on the isolate, for 48 hours to maintain humidity (95-100%). The plants were regularly sprayed with tap water. The experiment was repeated three times.

#### 2.3.4 Evaluation of parameters

#### 2.3.4.1 Incidence of attack

Young cashew plants showing symptoms of disease after contact were counted manually. The incidence was calculated using the following formula (Kranz, 1988):

$$I = (NPm / NP) X 100$$

With:

I: Incidence of attacking plants ; NPm: Number of diseased plants; NP: Total number of plants.

#### 2.3.4.2 Severity of attack

The severity of the disease on inoculated plants was determined using a visual rating scale

from 0 to 9 (Groth et al., 1999; Cardoso et al., 2004).

The severity index was calculated using the following formula from (Abu et al., 2013) :

$$Is = \sum (Xi * ni) / (N * Z)$$

With :

Is : Severity index ; Xi : Score assigned to the diseased plant ; ni : Number of diseased plants with the same score xi ; N : Total number of plants and Z : the highest score.

#### 2.3.4.3 Statistical analysis of the data

The data were analysed using the software (R 4.4.0). Following tests for normality and homogeneity of variance, the data were subjected to an analysis of variance. In the event of significant differences between the means, the Tukey's test was used to compare the means.

#### 3. RESULTS AND DISCUSSION

### 3.1. Results

#### 3.1.1 Symptoms induced by isolates on young cashew trees

The *Colletotrichum* isolates identified on the weeds induced two types of symptoms on the inoculated young cashew seedlings. These included black dots scattered across the leaf

blades and patches of greyish necrosis in the form of burning on the leaves. However, no disease symptoms were observed on the control plants (Fig. 2). Microbiological analysis of the infected leaves taken from the inoculated cashew trees enabled the initial fungi to be re-isolated.

#### 3.1.2 Incidence of symptoms

Fig. 3 shows the results relating to the incidence of symptoms induced by *Colletotrichum* sp. isolates on inoculated young cashew plants. The results showed that all inoculated cashew seedlings showed disease symptoms, in contrast to non-inoculated seedlings. Analysis of variance revealed a highly significant variation in incidence depending on the origin of the fungal isolates (P < 0.001). The highest incidence (90%) was obtained on plants inoculated with the CoIE2 isolate from the east. The lowest incidence (30%) was obtained on plants inoculated with the CoIN3 isolate from the north.

The results also showed a highly significant variation in the incidence of symptoms induced on young cashew seedlings by isolates from the same agroecological zone (P < 0.001). For isolates from the Centre agro-ecological zone, the highest incidence ((73.33%) was obtained on plants inoculated with the ColC1 isolate. The lowest incidence (50%) was found in plants inoculated with the ColC3 isolate. For isolates from the Centre-West agro-ecological zone, the highest incidence (43.33%) was obtained in plants inoculated with the ColC01 isolate and the lowest incidence (36.66%) was obtained in plants inoculated with the ColC02 isolate. For isolates from the eastern agro-ecological zone, the

highest incidence (90%) was obtained on plants inoculated with the CoIE2 isolate. The lowest incidence (56.66%) was obtained in plants inoculated with the CoIE1 isolate. For isolates from the Northern agro-ecological zone, the highest incidence (66.66%) was obtained on plants inoculated with the CoIN2 isolate. The lowest incidence (30%) was obtained in plants inoculated with the CoIN3 isolate.

#### 3.1.3 Symptom severity index

Fig. 4 shows the results of the evaluation of the symptom severitv index induced bv Colletotrichum sp. isolates on young cashew seedlings. The results showed that the incidence was high in the inoculated plants. Analysis of variance showed highly significant variation in symptom incidence according to the origin of the isolates (P < 0.001). In fact, the highest severity index of 7.66 was obtained in plants inoculated with the CoIN3 isolate from the North. The lowest severity index was 2.00 for plants inoculated with the CoIE3 isolate from the east.

The results also showed a highly significant variation in the severity index between isolates from the same agro-ecological zone (P < 0.001). In the Centre agro-ecological zone, the highest severity index (6.66) was obtained on plants inoculated with the ColC1 isolate. The lowest severity index (5) was obtained on plants inoculated with the ColC2 isolate. In the Centre-Ouest agro-ecological zone, the lowest severity index (3.66) was obtained on plants inoculated with the ColC01 isolate. And the lowest severity index (2.66) was obtained on plants inoculated with the ColC01 isolate. In the centre-Ouest severity index (2.66) was obtained on plants inoculated with the ColCO1 isolate. And the lowest severity index (2.66) was obtained on plants inoculated with the ColCO3 isolate. In the eastern agro-



Fig. 2. Symptoms induced on inoculated cashew plants (A): Control cashew plants; (B): Black dotted lines on inoculated cashew leaves and (C): Patches of necrosis on inoculated cashew leaves









ecological zone, the highest severity index (7.66) was obtained on plants inoculated with the ColE3 isolate and the lowest severity index (4.00) was obtained on plants inoculated with the ColE1 isolate. In the northern agro-ecological zone, the highest severity index (6.66) was obtained on plants inoculated with the ColN2 isolate and the lowest severity index (2.00) was obtained on plants inoculated with the ColN3 isolate.

# 3.2 Discussion

The results showed that all isolates induced symptoms characteristic of anthracnose in young cashew seedlings. This means that the isolates tested are pathogenic to cashew. Consequently, the weed hosts of these fungi would be alternative hosts of anthracnose in the cashew orchard. The symptoms identified on the weeds were black or grey dots scattered across the leaf blades and patches of reddish necrosis developing in the form of scorch marks on the weed leaves. These symptoms are similar to the anthracnose symptoms described on cashew in several previous studies (Afouda et al., 2013; Silué et al., 2017; Soro et al., 2022). These authors reported that Colletotrichum gloeosporioides is responsible for cashew anthracnose in Benin and Côte d'Ivoire. The twelve isolates tested belonged to the

gloeosporioides complex. These results corroborate those of (Muntala et al., 2020), who reported anthracnose symptoms on young cashew seedlings inoculated with a sporal suspension of Colletotrichum gloeosporioides. A difference in virulence was noted between the isolates. The ColE2 and ColE3 isolates present on V. paradoxa and A. zygia respectively in the eastern agro-ecological zone and the ColN1 isolate on D. oliveri in the northern agroecological zone were more virulent and aggressive than the others. The difference in aggressiveness observed could be explained by the fact that the isolates come from different species and belong to different agroecological zones. These results are similar to those of (Kouamé et al., 2011), who reported a difference in virulence between two isolates of Colletotrichum gloeosporioides inoculated into manages belonging to the keitt variety. These results could also be linked to the phenotypic characteristics of the isolates, which can vary according to the environmental conditions of the medium or the physicochemical composition of the host. This result is in agreement with those of (Photita et al., 2005; Martínez et al., 2009) who reported that a variation in the colour of the mycelium from white to grey or from pink to orange is possible and these variations are due to the host plant, the nature of the strain and the environmental conditions. The fungi re-isolated from infected cashew leaves are identical to the original isolates. Koch's postulate has therefore been verified. The fungi found to be pathogenic to cashew trees in this study came from D. oliveri, V. paradoxa, M. pruriens, B. sapida, A. zvaia, and P. erinaceus. These results attest to the presence of weed hosts of anthracnose in the cashew orchard. The presence of these weeds in the orchard would not be a problem if they were fully taken into account in anthracnose control strategies. But some of them, notably V. paradoxa and B. sapida, are deliberately kept in the orchards for their fruit, which is used for human consumption. Others are kept for their therapeutic properties (Sanders & Korsten, 2003). These plants thus escape cleaning operations and are not taken into account in fungicide treatments of orchards. The ability of fungi from these weeds to induce anthracnose in cashew trees remains a concern. The work of (Aboulaye et al., 2023) showed that D. oliveri, V. paradoxa, M. pruriens, B. sapida, A. zygia and P. erinaceus have high infection rates in ivorian cashew orchards. In addition to cashew, Colletotrichum gloeosporioides attacks other plants such as mango and avocado (Sanders &

Korsten. 2003: Pandev et al.. 2012). The ability of weeds to harbour fungal crop pathogens has been demonstrated in previous studies (Sales et al., 2012; Vasconcelos et al., 2012; De Oliveira et al., 2018; Alvarado-Huaman et al., 2019; Kumar et al., 2021). The work of (Alvarado-Huaman et al., 2019) showed that Cercospora coffeicola isolated from the weeds Cvathula achyranthoides (Kunth) and Chromoleana laevigata is responsible for iron spot of coffee. The same authors reported that Colletotrichum sp. isolated from Anthurium croatii is responsible for the anthracnose of coffee.

# 4. CONCLUSION

The isolates of Colletotrichum sp. present on weeds in the cashew orchard in Côte d'Ivoire are pathogenic to cashew. All twelve isolates tested induced symptoms in inoculated plants. The symptoms induced on young inoculated plants were typical of anthracnose. The weeds, D. oliveri, V. paradoxa, M. pruriens, B. sapida, A. zvgia, and P. erinaceus are alternative hosts of anthracnose in cashew orchards in Côte d'Ivoire. Their presence in the orchard certainly increases the risk of spreading anthracnose in the cashew some orchard. However, of them are permanently maintained in the orchard for their economic, nutritional, and therapeutic role. These trees should be regularly monitored in the same way as the cashew tree in order to reduce the risk of the disease spreading. For others, choosing a good cleaning period could help to control them better.

# DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declare that NO generative Al technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

# ACKNOWLEDGEMENTS

This work, which consisted in testing the pathogenicity of *Colletotrichum* sp isolates from weeds on cashew trees, is the first of its kind in Côte d'Ivoire. It was carried out with the financial and material support of the Fonds Interprofessionnel pour la Recherche et le Conseil Agricoles (FIRCA), the Projet de Promotion de la Compétitivité de la Chaîne de valeur de l'Anacarde (PPCA), the Conseil du

Coton et de l'Anacarde (CCA), the Programme National de Recherches sur l'Anacarde (PNRA).

#### **COMPETING INTEREST**

Authors have declared that no competing interests exist.

# REFERENCES

- Aboulaye, T., Sibirina, S., Emmanuella, A. A. R., Karidia, T.-O., Abo, K., & Daouda, K. (2023). Fungal flora on weeds in the cashew (*Anacardium occidentale* L.) orchard in Côte d'Ivoire. *American Journal* of *Plant Sciences*, 14, 448–463. https://doi.org/10.4236/ajps.2023.144029
- Abu, B. A., Nur Ain, I. M., & Umi, K. Y. (2013). Diversity of *Fusarium* species associated with post-harvest fruit rot disease of tomato. *Sains Malaysiana*, *4*2, 911– 920.
- Afouda, L. C. A., Zinsou, V., Balogoun, R. K., Onzo, A., & Ahohuendo, B. C. (2013). Inventaire des agents pathogènes de l'anacardier (*Anacardium occidentale* L.) au Bénin. *Bulletin de Recherche Agronomique du Bénin, 73*, 13–19.
- Agostini-Costa, T. S., Faria, J. P., Naves, R. V., & Vieira, R. F. (2006). Cajus do cerrado. In R. F. Vieira, T. S. Agostini-Costa, D. B. Silva, F. R. Ferreira, & S. M. Sano (Eds.), *Frutas Nativas da Região Centro-Oeste do Brasil* (1st ed., pp. 136–151). Embrapa Recursos Genéticos e Biotecnologia.
- Alvarado-Huaman, L., Castro-Cepero, V., Tejada-Soraluz, J. L., Borjas-Ventura, R., & Julca-Otiniano, A. (2019). Fungi and nematodes associated with weeds in the coffee crop (*Coffea arabica* L.) in the Central jungle of Perú. *Revista de Investigación e0020vn: Innovación Agropecuaria y de Recursos Naturales*, 6(2), 37–45.
- Cardoso, J. E., Santos, A. A., Rossetti, A. G., & Vidal, J. C. (2004). Relationship between incidence and severity of cashew gummosis in semiarid northeastern Brazil. *Plant Pathology*, 53, 363–367. https://doi.org/10.1111/j.0032-0862.2004.01007.x
- De Oliveira, E. F., Dos Santos, P. R. R., & Dos Santos, G. R. (2018). Seeds of weeds as an alternative host of phytopathogens. *Arquivos do Instituto Biológico, 85*, 1–7, e0972017.

- Dentika, P., Ozier-Lafontaine, H., & Penet, L. (2021). Weeds as pathogen hosts and disease risk for crops in the wake of a reduced use of herbicides: Evidence from yam (*Dioscorea alata*) fields and *Colletotrichum* pathogens in the tropics. *Journal of Fungi*, 7, 283.
- Figueiredo, L. C., Figueirêdo, G. S., Quecine, M. C., Cavalcanti, F. C. N., Santos, A. C., & Costa, A. F. (2012). Genetic and pathogenic diversity of *Colletotrichum gloeosporioides*, the causal agent of cashew anthracnose. *Indian Journal of Fundamental and Applied Life Sciences*, 2, 250–259.
- Groth, J. V., Ozmon, E. A., & Busch, R. H. (1999). Repeatability and relationship of incidence and severity measures of scab of wheat caused by *Fusarium graminearum* in inoculated nurseries. *Plant Disease*, *83*, 1033–1038.
- Houndahouan, D. E. T., Zanou, A., Sikirou, R., Adomou, A., Zinsou, V., Boukari, S., & N'djolossè, K. (2018). Les pertes économiques dues à l'anthracnose de l'anacardier au Bénin. *European Scientific Journal, 14*(15), 127–138. https://doi.org/10.19044/esj.2018.v14n15p 127
- Kouadio, K. F. (2015). Contributions des biotechnologies à la sécurité alimentaire : cas du biofertilisant organique symbiose (Anabaena-Azollae et Azolla filiculoides) sur oryza sativa (riz CB-one) en Côte d'Ivoire (Master's thesis, UFR Agroforesterie, Université Jean Lorougnon Guédé de Daloa, 50p).
- Kouamé, G. K., Sorho, F., Koné, D., Bomisso, L. E., Aké, S., & Yatty, J. (2011). Activité pathologique comparée de deux isolats de *Colletotrichum gloeosporioides* (PENZ.) sur deux variétés de mangues (*Mangifera indica* L.). Agronomie Africaine, 23, 33–41.
- Kranz, J. (1988). Measuring plant disease. In J. Kranz & J. Rotem (Eds.), *Experimental techniques in plant disease epidemiology* (pp. 35–50). Springer.
- Kumar, S., Bhowmick, M. K., & Ray, P. (2021). Weeds as alternate and alternative hosts of crop pests. *Indian Journal of Weed Science*, *53*(1), 14–29.
- Martínez, E. P., Hío, J. C., Osorio, J. A., & Torres, M. F. (2009). Identification of *Colletotrichum* species causing anthracnose on Tahiti lime, tree tomato,

and mango. *Agronomía Colombiana,* 27, 211–218.

- Muntala, A., Norshie, P. M., Santo, K. G., & Saba, C. K. S. (2020). *Colletotrichum gloeosporioides* species complex: Pathogen causing anthracnose, gummosis, and die-back diseases of cashew (*Anacardium occidentale* L.) in Ghana. *European Journal of Agriculture and Food Sciences*, 2(6), 1–10. https://doi.org/10.24018/ejfood.2020.2.6.14 6
- Pandey, A., Yadava, L. P., Mishra, R. K., Pandey, B. K., Muthukumar, M., & Chauhan, U. K. (2012). Study on the pathogenesis incident and of Colletotrichum gloeosporioides Penz., cause of anthracnose of mango. International Journal of Sciences and Nature, 3, 220-232.
- Photita, W., Taylor, P. W., Ford, R., Hyde, K. D., & Lumyong, S. (2005). Morphological and molecular characterization of *Colletotrichum* species from herbaceous plants in Thailand. *Fungal Diversity*, 18, 117–133.
- RGPH (2021). Recensement général de la population et de l'habitat en Côte d'Ivoire.
- Sales, J. R., Oliveira, O. F., Medeiros, E. V., Guimarães, I. M., Correia, K. C., & Michereff, S. J. (2012). Ervas daninhas como hospedeiras alternativas de patógenos causadores do colapso do meloeiro. *Revista Ciência Agronômica*, 43(1), 195–198.
- Sanders, G. M., & Korsten, L. (2003). A comparative morphology of South African avocado and mango isolates of *Colletotrichum gloeosporioides*. *Canadian Journal of Botany, 81*, 877– 885.
- Silué, N., Abo, K., Johnson, F., Camara, B., Koné, M., & Koné, D. (2018). Evaluation in vitro et in vivo de trois fongicides de synthèse et d'un fongicide biologique sur la croissance et la sévérité de Colletotrichum gloeosporioides et de Pestalotia heterornis, champignons responsables de maladies foliaires de l'anacardier (Anacardium occidentale L.) en Côte d'Ivoire. Agronomie Africaine, 30(1), 107–122.
- Silué, N., Soro, S., Koné, T., Abo, K., Koné, M., & Koné, D. (2017). Parasitical fungi in cashew (*Anacardium occidentale* L.)

orchard of Côte d'Ivoire. *Plant Pathology Journal, 16,* 82–88.

Soro, A. N., Soro, S., Yéo, G., Kouman, A. M. N., Tehua, A. A., Silué, N., Abo, K., & Koné, D. (2022). Severity and distribution updated of anthracnose on cashew trees (*Anacardium occidentale* L.) in Côte d'Ivoire. *Agronomie Africaine Sp., 34*(1), 49–58. https://www.ajol.info/index.php/aga/article/

view/229507

- Soro, S., N'Da, A. A., & Koné, D. (2013). Comportement des génotypes d'anacardiers (*Anacardium occidentale* L.) à l'anthracnose (*Colletotrichum gloeosporioides* Penz.) au Nord de la Côte d'Ivoire. (AGRAR), Yamoussoukro, Côte d'Ivoire, 19p.
- Soro, S., Silué, N., Ouattara, G. M., Chérif, M., Camara, I., Sorho, F., Ouali, N. M., Abo, K., Koné, M., & Koné, D. (2015). Investigations on major cashew diseases in Côte d'Ivoire. *Proceedings of the Third International Cashew Conference*, 16-19 November 2015, Dar es Salaam, Tanzania, 158–166.
- Vasconcelos, M. C. C., Silva, A. F. A., & Lima, R. S. (2012). Interferência de plantas daninhas sobre plantas cultivadas. *Agropecuária Científica no Semi-Árido,* 8(1), 1–6.
- Veloso, J. S., Câmara, M. P. S., Lima, W. G., Michereff, S. J., & Doyle, V. P. (2018). Why species delimitation matters for fungal ecology: *Colletotrichum* diversity on wild and cultivated cashew in Brazil. *Fungal Biology*, 122(7), 677–691. https://doi.org/10.1016/J.FUNBIO.2018.03. 005
- Veloso, J. S., Duarte, I. G., Farias, O. R. D., & Câmara, M. P. S. (2022). Antracnose do cajueiro: Etiologia, sintomatologia e aspectos epidemiológicos. *Revisão Anual de Patologia de Plantas, 28*(July), 200– 215. https://doi.org/10.31976/0104-038321v280009
- Wonni, I., Sereme, D., Ouedraogo, I., Kassankagno, A. I., Dao, I., Ouedraogo, L., & Nacro, S. (2017). Diseases of cashew nut plants (*Anacardium* occidentale L.) in Burkina Faso. Advances in Plants & Agriculture Research, 6(3), 78– 83. https://doi.org/10.15406/apar.2017.06.002

16

#### Zro, B. G. F., Guéi, A. M., Nangah, K. Y., Soro, D., & Bakayoko, S. (2016). Statistical approach to the analysis of the variability

and fertility of vegetable soils of Daloa (Côte d'Ivoire). *African Journal of Soil Science*, *4*(4), 328–338.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/128488