

International Journal of Plant & Soil Science

Volume 37, Issue 1, Page 67-75, 2025; Article no.IJPSS.118982 ISSN: 2320-7035

# Genetic Variability and Genetic Advance Studies for Bulb Yield and Its Components of Different Mutant Lines of Garlic (*Allium sativum* L.)

# Ayusha Gupta <sup>a\*</sup> and Pravin Kumar Sharma <sup>a</sup>

<sup>a</sup> Department of Vegetable Science, COA, Indira Gandhi Krishi Vishwavidyalaya, Raipur (C.G), India.

#### Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

#### Article Information

DOI: https://doi.org/10.9734/ijpss/2025/v37i15252

#### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/118982

**Original Research Article** 

Received: 17/10/2024 Accepted: 21/12/2024 Published: 10/01/2025

# ABSTRACT

This study aims to study aims to evaluate the genetic variability and genetic advance for bulb yield and components of different mutant lines of Garlic (*Allium sativum L.*) under Chhattisgarh plains. The experiment was laid out in Randomized Block design with three replications 21 mutant lines *viz.*, IG C-1, IG C-2, IG C-3, local check, IG M-2021-1, IG M-2021-2, IG M-2021-3, IG M-2021-4, IG M-2021-5, IG M-2021-6, IG M-2021-7, IG M-2021-8, IG M-2021-9, IG M-2021-10, IG M-2021-11, IG M-2021-12, IG M-2021-13, IG M-2021-14, IG M-2021-15, IG M-2021-16, IG M-2021-17. Data were analyzed to work out performance of mutant lines, to assess genetic variability, heritability, genetic advance. The character taken under study include plant height (cm), number of leaves, collar height (cm), collar thickness (cm), fourth leaf length (cm), fourth leaf width (cm), polar diameter (cm), equatorial diameter (cm), neck thickness (cm), number of cloves, average weight of

<sup>\*</sup>Corresponding author: E-mail: ayushagupta5224@gmail.com;

*Cite as:* Gupta, Ayusha, and Pravin Kumar Sharma. 2025. "Genetic Variability and Genetic Advance Studies for Bulb Yield and Its Components of Different Mutant Lines of Garlic (Allium Sativum L.)". International Journal of Plant & Soil Science 37 (1):67-75. https://doi.org/10.9734/ijpss/2025/v37i15252.

bulb (g), TSS, days to maturity, weight of ten outer cloves (g), marketable yield (t ha<sup>-1</sup>), total yield (t ha<sup>-1</sup>). Estimates of genetic parameters for various character under study revealed that high genotypic co-efficient of variation was found in general compared to phenotypic co-efficient of variation. Heritability estimates under present investigation ranged from moderate to high. High heritability estimates recorded for collar height, weight of outer ten cloves, marketable yield, total yield. High heritability coupled with high genetic advance was recorded for collar height, marketable yield and bulb yield.

Keywords: Heritability; genetic advance; mutant lines etc.

#### **1. INTRODUCTION**

Garlic (Allium sativum L.) is an annual aromatic herbaceous spice that belongs to the Alliaceae family (Kurian, 1995). After onion, it is the second most widely cultivated spice crop (Rubatzky, and Yamaguchi, 1997). Garlic dates back to prehistoric times in Central Asia and the Mediterranean region (Thompson and Kelly, 1957). Before 2000 B.C., garlic cultivation was discovered in Egypt, and nearly 5000 years ago in India (Kamenetsky and Rabinowitch, 2001). Garlic is produced worldwide. It is consumed as and also spice. flavouring agent has pharmaceutical benefits. "Allium" is the widest and the most diverse representative genus of the Alliaceae. It is spread across in North Africa, North America, Europe and Asia and contains around 700 different species. China is the leading producer of garlic and accounts for 20.7 million tons of production, followed by India with production of 2.9 million. In India, Madhya Pradesh is the leading producer of followed by other states like Rajasthan, Uttar Pradesh, and Gujarat (Anon., 2021a). Whereas in Chhattisgarh it occupies 4,411 ha area with production of around 26,185 MT (Anon., 2021b). In recent years, mutation breeding has gained popularity as a convenient tool for crop development and as a way to complement current germplasm for cultivar improvement in breeding programmes (Acharya, et al., 2007). Furthermore, mutation are defined as a sudden heritable changes in gene. It can be either spontaneous or induced. It can be induced either in seed or propagating material.(Al safadi et al., 2000) (

#### 2. MATERIALS AND METHODS

The study was conducted in Rabi season 2021-2022 at Research cum Demonstration Farm, College of Agriculture, Indira Gandhi Krishi Vishwavidyalaya, Raipur (C.G.) - 492012. It lies between 21°16'N latitude and 81°36'E longitude at an altitude of 289.56 meters above mean sea level.

The phenotypic and genotypic variance components, and coefficients of phenotypic and genotypic coefficients of variation were calculated by the methods suggested by Burton and Devane, 1953 is illustrated below:

1) Genotypic coefficient of variation (GCV) -

$$\mathsf{GCV}(\%) = \frac{\sqrt{\sigma^2} g}{\bar{X}} \times 100$$

Where,

 $\sigma^2$  g = genotypic variance

 $\overline{X}$  = Mean of characters

2) Phenotypic coefficient of variation (PCV) -

PCV (%) = 
$$\frac{\sqrt{\sigma^2} p}{\bar{X}} \times 100$$

Where,

 $\sigma^2 p$  = phenotypic variance

 $\overline{X}$  = Mean of characters

#### 2.1 Heritability

Heritability with reference to broad sense (h<sup>2</sup>b) is defines as proportion of genotypic variance to total variance (phenotypic variance). It was estimated by using Hansen, et al., (1956) formula:

$$h^2 b = \frac{\sigma^2 g}{\sigma^2 p} \times 100$$

The heritability was classified into low, moderate and high as suggested by Robinson, (1966):

<50%	=	Low		
50-70% =	Mode	Moderate		
>70%	=	High		

### 2.2 Genetic Advance

It was calculated as per method given by Johnson, et al., (1955)

 $GA = k \sigma ph^2$ 

Where,

k = Selection intensity at 5% i.e. 2.06 (constant)

 $\sigma p$  = Phenotypic standard deviation

 $h^2$  = Heritability in broad sense

Genetic advance was classified into high, moderate and low as suggested by Johnson, et al. (1955) > 20% = High

10-20% = Moderate

#### 2.3 Genetic Advance as Percentage of Mean

It was estimated by the formula -

**GA (%)** =  $\frac{GA}{X} \times 100$ 

Where, GA = Genetic advance

 $\overline{X}$  = Mean of character

#### 3. PARAMETERS UNDER STUDY

The estimates of PCV and GCV were classified as low (<10%), moderate (10-12%) and high (>20%) according to Sivasubramanian and Madhavamenon, (1973). Data presented in Table 1 indicated highest range of variation (17.73-42.94) for fourth leaf width followed by plant height (14.55-58.75). For neck thickness the range of variation was very narrow (0.36-1.18). The Genotypic coefficient variation (GCV) and Phenotypic coefficient variation (PCV) garlic mutant lines are depicted in Fig. 1 and Fig. 2. High magnitude of genotypic as well as phenotypic coefficient of variation were recorded for traits viz., Collar height (33.41 and 35.00), Weight of ten outer cloves (33.26 and 39.84), Marketable yield (32.31 and 33.77), total yield (26.22 and 28.01) and average weight of bulb (19.18 and 23.31) suggests the substantial improvement can be achieved through selection of these traits. Moderate PCV and GCV was recorded for fourth leaf length (16.12 and 17.28), this suggest existence of considerable variability in the population. Selection based on these traits will contribute in improvement in genotype. Remaining character had low PCV and GCV. Lowest PCV and GCV recorded for number of leaves (5.37 and 3.11). The result of the present study were found to be close to the findings of Islam, et al. (2004), Zahedi, et al. (2007), Tsega, et al. (2010), Singh, et al. (2011), Kumar, et al. (2015), Vatsayan, et al. (2015) and Sharma, et al. (2016). They recorded high PCV and GCV for Plant height, average weight of bulb, number of cloves per bulb and marketable yield.

The genetic advance as percentage of mean ranged from 3.20 to 65.70. Highest genetic advance as percentage of mean was recorded Collar height (65.70%) followed for by marketable yield (63.68%), weight of ten outer cloves (57.19%) total yield (50.58%). The genetic advance as percentage of mean was found to moderate to high for all characters except number of leaves (3.72%), fourth leaf width (8.37%), days to maturity (3.20%). The high value of genetic advance for the traits under study revealed that these characters are governed by additive genes and selection will be rewarding for further improvements in future. Moderate genetic advance for traits suggesths that both additive and non-additive variance are operating in these traits. Furthermore the traits low genetic advance showina indicates significance of non-additive gene effects. The genetic advance as percentage of mean is presented in Table 2 and Fig. 2. The findings are in close associations with reports of Tsega, et al. (2010), Sonkiya, et al. (2012), Singh, et al. (2011), Sable, et al. (2020), Anand (2022), Beena et al. (2021), Kar (2013).

#### 4. RESULTS AND DISCUSSION

Heritability estimates for most of the character ranged between moderate to high except number of leaves and fourth leaf width (cm). High heritability coupled with high genetic advance were recorded for collar height (cm), marketable yield (t ha-1) and total yield (t ha-1) whereas moderate level of heritability with moderate level of genetic advance was observed for fourth leaf length (cm), neck thickness (cm), average weight of bulb (g). The result of the present study were found to be close to the findings of Tsega, et al., (2010), studied about 25 indigenous garlic accessions to determine variabilitv and correlation for yield traits and bulb yield. Little difference was observed between phenotypic and genotypic coefficient of variation which indicates that variability in accessions was

S.No.	Parameters	Range		Mean	Coefficient of variation (%)		h² (b)	Genetic advance	G.A as % of mean
	Characters	Minimum	Maximum		Genotypic	Phenotypic			
1	Plant height (cm)	41.55	58.75	51.723	10.109	13.174	58.883	8.265	15.980
2	Number of leaves	8.3	9.27	8.692	3.116	5.375	33.593	0.323	3.720
3	Collar height (cm)	5.4	16.38	8.686	33.417	35.009	91.111	5.708	65.707
4	Collar thickness (cm)	2.59	3.96	2.965	8.463	10.290	67.651	0.425	14.340
5	Fourth leaf length (cm)	17.73	42.94	33.945	16.126	17.286	87.029	10.519	30.990
6	Fourth leaf width (cm)	0.58	1.40	1.217	9.355	21.529	18.882	0102	8.374
7	Polar diameter (cm)	2.63	3.49	3.197	7.387	9.063	66.426	0.396	12.402
8	Equitorial diameter (cm)	2.56	3.93	3.336	9.284	10.902	72.528	0.543	16.288
9	Neck thickness (cm)	0.36	1.18	0.699	22.149	29.998	54.515	0.235	33.688
10	Weight of 10 outer cloves (g)	3.77	11.43	7.394	33.261	39.849	69.668	4.229	57.190
11	Number of cloves	19.47	28.33	23.670	9.241	10.314	80.272	4.037	17.055
12	Average bulb weight (g)	9.27	17.95	13.446	19.189	23.315	67.737	4.374	32.533
13	TSS	32.26	40.22	36.286	6.753	7.452	82.131	4.575	12.608
14	Days to maturity	130.33	141.33	136.397	2.043	2.687	57.807	4.364	3.199
15	Maretable yield (t/ha)	2.88	10.04	6.513	32.31	33.774	91.528	4.148	63.68
16	Total yield (t/ha)	3.61	10.31	7.170	26.228	28.015	87.649	3.627	50.584

# Table 1. Genetic parameter of variation for bulb yield and its components in garlic

S.No.	Characters	GCV %	PCV (%)	h2(b) (%)	Genetic advance as % of mean
1	Plant height (cm)	М	М	Μ	L
2	Number of leaves	L	L	L	L
3	Collar height (cm)	Н	Н	Н	Н
4	Collar thicness (cm)	L	Μ	Μ	L
5	Fourth leaf length (cm)	Μ	Μ	Н	Μ
6	Fourth leaf width (cm)	L	Н	L	L
7	Polar diameter (cm)	L	L	Μ	L
8	Equitorial diameter (cm)	L	Μ	Н	L
9	Neck thicness (cm)	Н	Н	Μ	Μ
10	Weight of 10 outer cloves (g)	Н	Н	Μ	Н
11	Number of cloves	L	Μ	Н	L
12	Average bulb weight (g)	Μ	Н	Μ	Μ
13	TSS	L	L	Н	L
14	Days to maturity	L	L	Μ	L
15	Maretable yield (t/ha)	Н	Н	Н	Н
16	Total yield (t/ha)	Н	Н	Н	Н

 Table 2. Summary of Genotypic coefficient of variation (GCV), Phenotypic coefficient of variation (PCV), heritability (h2b) and genetic advance as percent of mean of bulb yield and its components in garlic

H- HIGH; M- MEDIUM; L-LOW





Fig. 1. Graphical representation of GCV of bulb yield and its component in garlic



Fig. 2. Graphical representation of GCV of bulb yield and its component in garlic

characterized by their genetic constitution, Sonikiya, et al., (2012), studied about 10 genotypes for 14 quantitative traits in garlic and result depicted high genetic advance and heritability. high genetic advance was reported as percent of mean for weight of 100 cloves per plant and leaf area, Kumar, et al., (2015) studied about 41 genotypes of garlic under Jammu region for quality, yield and other agronomic characters. It was found that the genotypes showed high heritability along with high genetic advance as a percent of mean as observed for average weight of cloves, yield per hectare, plant heigh, leaf length, average weight of bulb and equatorial diameter. The analysis of various result showed that genetic variability of all traits except for traits like dry matter content, total soluble solids, leaves per plant, leaf length, and equatorial diameter. Sharma, et al., (2016) evaluated 131 diverse genotypes of garlic. High heritability was observed for all traits under study coupled with moderate level of genetic advance for characters like bulb weight per plant, clove weight, polar diameter of bulb equatorial diameter of bulb, pseudostem height, and number of cloves per bulb. High estimates of PCV and GCV were observed for bulb weight per plant, plant height, number of cloves per bulb. pseudo stem and leaf length. Selection of these traits would be rewarding towards higher bulb yield. Occurrence of high heritability coupled with high genetic advance for marketable yield in tonnes per hectare, total yield of bulb in tonnes per height hectare. collar indicating preponderance of additive genetic variance in the genetic control of these traits. Selection would be more effective for these characters towards the improvement of total bulb yield in garlic.

Heritability ranged from 18.88-91.52%. Highest estimate were recorded for marketable yield (91.52%), collar height (91.11%), total yield (87.64%), number of cloves (80.27%). High heritability for above characters represented that they were least affected by environmental fluctuation and selection based on phenotypic expression may be reliable. The finding are similar to findings of Tsega et al. (2010),Tesfaye et al. (2021), Kar (2013), Futane (2000), Bhatt et al., (2017) Meena et al., (2020), Tripathi e al (2013).I

# 5. CONCLUSION

Mutation breeding has gained popularity as a convenient tool for crop development and as a way to complement current germplasm for cultivar improvement in breeding programmes. High heritability estimates recorded for collar height, weight of outer ten cloves, marketable yield, total yield. High heritability coupled with high genetic advance was recorded for collar height, marketable yield and bulb yield.

# DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative Al technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

# COMPETING INTERESTS

Authors have declared that no competing interests exist.

# REFERENCES

Acharya, S.N. et al. (2007) Improvement in the medicinal and nutritional properties of

fenugreek (Trigonella foenum-graecum L.). In: Acharya, S.J. and Thomas, J.E. (eds) Advances in Medicinal Plant Research. Research Signpost, Kerala, India, pp. 491– 512.

- Al-Safadi, B., Mir Ali, N., & Arabi, M. I. E. (2000). Improvement of garlic (*Allium sativum* L.) resistant to white rot and storability using γ-irradiation induced mutations. *Journal of Genetics and Breeding, 54*(3), 175-181.
- Anand, M., Sankari, A., Kamalkumaran, P. R., & Velmurugan, M. (2022). Studies on the genetic variability, character association and path coefficient analysis in garlic (*Allium sativum* L.). *Electronic Journal of Plant Breeding, 13*(1), 69-74.
- Anonymous. (2021a). Production volume of garlic in India FY 2022. *Statista Research Department*.
- Anonymous. (2021b). Directorate Horticulture and Farm Forestry, Chhattisgarh (Department of Agriculture, Government of Chhattisgarh).
- Beena, R., Veena, V., Jaslam, M. P., Nithya, N., & Adarsh, V. S. (2021). Germplasm innovation for high-temperature tolerance from traditional rice accessions of Kerala using genetic variability, genetic advance, path coefficient analysis and principal component analysis. *Journal of Crop Science and Biotechnology, 24*(5), 555-566.
- Bhatt, R., Soni, A. K., Jangid, K., & Kumar, S. (2017). A study on genetic variability, character association and path analysis in promising indigenous genotypes of garlic (*Allium sativum* L.). *International Journal of Pure and Applied Biosciences, 5*(1), 679-687.
- El-Fiki, A., & Adly, M. (2019). Morphological, molecular, and organosulphur compounds characterization in irradiated garlic (*Allium sativum*) by GC–MS and SCoT markers. *Journal of Radiation Research and Applied Sciences, 13*(1), 61–70.
- Futane, N. W., Jogdande, N. D., Gonge, V. S., Warade, A. D., & Khandagale, S. S. (2006). Evaluation of garlic (*Allium sativum* L.) genotypes. *International Journal of Agriculture Science*, 1, 4-5.
- Hanson, C.H., H.F. Robinson and R.E. Comstock. 1956. Biometrical studies on yield in segregating population of Korean lespedesa. Agron. J. 48:268-272.

- Islam, M. J., Islam, M. A., Aktertania, S., Saha, S. R., Alam, M. S. & Hasan, M. K. (2004). Performance of evaluation of some garlic genotypes in Bangladesh. Asian J. Plant Sci., 3(1): 14-16.
- Johnson, W.W., H.F. Robinson and R.E. Comstock. 1955. Genotypic and phenotypic correlations in soybeans and their implications in selection. Agron. J. 47:477-482.
- Kar, S. (2013). Study on garlic (*Allium sativum* L.) genotypes under Chhattisgarh plains.
   M.Sc. (Ag.) Thesis, Indira Gandhi Krishi Vishwavidyalaya, Raipur, 7-15.
- Kamenetsky and Rabinowitch, 2001. Flower development in bolting garlic. Sexual plant reproduce. 4:235-241.
- Kumar, G. P., Sunil, N., Sekhar, J. C., & Srinivasa Chary, D. (2024). Assessment of genetic variability, heritability and genetic advance in maize genotypes (*Zea mays* L.). *Journal* of Experimental Agriculture International, 46(3), 146-155. https://doi.org/10.9734/jeai/2024/v46i3233 3
- Kurian JC (1995). Plant that Heal (1st edn.), Oriental Watchman publishing House, Punj., India. P31.
- Meena, M. L., Yadav, S., Kumar, S., Singh, R., Abha, R., & Mandel, R. K. (2020). Genetic variability, heritability and genetic advances in the garlic, *Allium sativum* genotypes. *Bioscience Biotechnology Research Communication, 13*(1).
- Robinson HF (1966). Quantitative genetics in relation to breeding of the centennial of mendalism. Indian J Genet. 26: 171-187.
- Rubatzky VE, Yamaguchi M (1997). World Vegetables. Principles, production and nutritive values. 2nd ed. Chapman and Hall. International Thomson publishing, New York, USA.
- Sable SV, Deshmukh DT, Ghawade SM, Rawat SS (2020) Genetic Variability and Correlation Studies in Garlic (*Allium sativum* L.). Int. J. Curr. Micro biol. App. Sci, (2020) 9(5): 358-363.
- Sharma, R., Omotayo, K., Kattula, N., Malik, S., Kumar, M., & Sirohi, A. (2016). Genetic variability, heritability and genetic advance in garlic genotypes. *International Journal of Agriculture Sciences, 24*(8), 2894-2898.
- Singh SR, Mir JI, Ahmed N, Rashid R, Wani S, Sheikh MA, Shafi W, Jan N and Mir H

(2011). RAPD profile based grouping of garlic germplasm with respect to photoperiodism.*J. Tropical Agric.* 49(1-2): 114-117.

- Sivasubramanian J and Madhavamenon P 1973. Genotypic and phenotypic variability in rice. Madras Agric. J. 12: 15-16.
- Sonkiya ,A.K.,Singh ,P.P. and Naruka, I.S. 2012. Variability, character association and path coefficient analysis in garlic. Medicinal Plant .4(2) : 90-93
- Tesfaye, A., Mijena, D. F., Zeleke, H., & Tabor, G. (2021). Genetic variability and character association for bulb yield and yield related traits in garlic in Ethiopia. *African Crop Science Journal, 29*(2), 293-308.
- Thompson, H.C. and Kelly, W.C. (1957) Vegetable Crops. McGraw Hill Book Company Inc., New York.
- Tripathi, A., Mishra, D. K., & Shukla, J. K. (2013). Genetic variability, heritability and genetic advance of growth and yield components of Jatropha (*Jatropha curcas* Linn.) genotypes. *Trees*, *27*, 1049-1060.
- Zahedi B, Kashi AK, Zamani Z, Mosahebi GH and Hassani M (2007). Evaluation of Iranian Garlic Allium sativum Lgenotypes using Multivariate analysis method based on Morphological characteristics. Biotechnology. 63: 353-356
- Zahedi B, Kashi AK, Zamani Z, Mosahebi GH and Hassani M (2007). Evaluation of Iranian Garlic Allium sativum Lgenotypes using Multivariate analysis method based on Morphological characteristics. Biotechnology. 63: 353-356
- Tsega, K., Tiwari, A., & Woldetsadik, K. (2010). Genetic variability, correlation and path coefficient among bulb yield and yield traits in Ethiopian garlic germplasm. *Indian Journal of Horticulture*, 67(4), 489-499.
- Zahedi B, Kashi AK, Zamani Z, Mosahebi GH and Hassani M (2007). Evaluation of Iranian Garlic Allium sativum Lgenotypes using Multivariate analysis method based on Morphological characteristics. Biotechnology. 63: 353-356
- Zahedi B, Kashi AK, Zamani Z, Mosahebi GH and Hassani M (2007). Evaluation of Iranian Garlic Allium sativum Lgenotypes using Multivariate analysis method based on Morphological characteristics. Biotechnology. 63: 353-356

#### Zahedi B, Kashi AK, Zamani Z, Mosahebi GH and Hassani M (2007). Evaluation of Iranian Garlic *Allium sativum*Lgenotypes

using Multivariate analysis method based on Morphological characteristics. *Biotechnology.* 63: 353-356.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/118982